

Availability Authorization (DAA)

The DAA approval process applies to all forms of published NASA Scientific and Technical Inform the responsible NASA Project Officer, Technical Monitor, author, or other appropriate NASA offici Explanations are on the back of this form and are presented in greater detail in NPG 2200.2, "Gu Technical Information."	nation (STI), whether disseminated in print or electronically. It is to be initiated by iaid for all presentations, reports, papers, and proceedings that contain NASA STI. Original idelines for Documentation, Approval, and Dissemination of NASA Scientific and Modified		
I. DOCUMENT/PRO.	JECT IDENTIFICATION		
Estimates of plant canopy attenuation, absorption and reflectance signatures and their relationships to select biophysical and biochemical features as determined by Optimal Passive Ambient Correlation Spectroscopy (OPACS).	AUTHOR(S) Hall, C.R., C.R. Bostater, and M.F. Pegaz. Pp. 527-532. In: Proceedings of Eco-Informa'96: Global Networks for Environmental Information, Volume 10. Environmental Research Institute of Michigan, Ann Arbor.		
	1		
ORIGINATING NASA ORGANIZATION	PERFORMING ORGANIZATION (If different)		
Kennedy Space Center Biomedical Office	Dynamac Corporation, KSC		
CONTRACT/GRANT/INTERAGENCY/PROJECT NUMBER(S) DOCUM	ENT NUMBER(S) DOCUMENT DATE		
See A	Authors Block. 1996		
For presentations, documents, or other STI to be externally published (including through electronic media), enter appropriate information on the intended publication such as name, place, and date of conference, periodical, or journal name, or book title and publisher in the next box. These documents must be routed to the NASA Headquarters or Center Export Control Administrator for approval (see Sections III and VIII).			
CHECK ONE (One of the five boxes denoting Security Classification must	be checked.)		
SECRET SECRET RD CONFIDE	ENTIAL CONFIDENTIAL RD X UNCLASSIFIED		
AVAILABIL.	ITY CATEGORY AS EAST OF THE PARTY OF THE PAR		
ITAR EAR Export Controlled Document - USML Category /CCL Export Control Classification Number (ECCN) (Documents marked in this block must have the concurrence/approval of the NASA Headquarters or Center Export Control Administrator (see Section VIII).)			
Confidential Commercial Document (check appropriate box at left and indicate below the appropriate limitation and expiration): TRADE SECRET U.S. Government agencies and U.S. Government agency contractors only NASA contractors and U.S. Government only U.S. Government agencies only NASA personnel and NASA contractors only NASA personnel only Available only with the approval of issuing office: Limited until (date)			
AVAILABLE confidential commercial data, and should	nclassified, may not be export controlled, may not contain trade secret or d have cleared any applicable patents application process. OSING AN INVENTION		
	OR INTELLECTUAL PROPERTY COUNSEL SIGNATURE DATE		
V. BLANKET RELI	EASE (OPTIONAL)		
All documents issued under the following contract/grant/project numay be processed as checked in Sections II and III. The blanket release authorization granted on (date) is RESCINDED - Future documents must have individual available in MODIFIED - Limitations for all documents processed in the blocks as checked in Sections II and III.	ē e		

	lential commerc	ial information, and/or discloses an invention for which a patent has	s been applied, and	
the appropriate limitation is checked in does NOT contain export controlled, or applied, and may be released as indica	onfidential comm	d/or IV. nercial information, nor does it disclose an invention for which a pat	ent has been	
NAME OF AUTHOR/ORIGINATOR Carlton Hall	MAIL CODE Dyn-2	SIGNATURE Hall	DATE 4 6 98	
VII. PROJE	CT OFFICER/T	ECHNICAL MONITOR/DIVISION CHIEF REVIEW	1	
APPROVED FOR DISTRIBUTION AS MARK				
NAME OF PROJECT OFFICER OR TECH. MONITOR RAYMONID M. WHEELER	MAIL CODE JJ-G	SIGNATURE M Weller	DATE 4/8/98	
VIII. EXPORT CONTROL REVIEW/CONFIRMATION				
Public release is approved Export controlled limitation is approved	/	ort controlled limitation is not applicable ort controlled limitation (ITAR/EAR) marked in Section III is assigned to this do	ocument:	
USML CATEGORY NUMBER CCL ECCN NUM	BER	HO OR CENTER EXPORT CONTROL ADMINISTRATOR (as applicable)	DATE 4/16/98	
IX. PROGRAM OFFICE OR DELEGATED AUTHORITY REVIEW				
APPROVED FOR DISTRIBUTION AS MARI	CED ON REVERS	NOT APPROVED CODE M, DAA	LETTER,6/14/	
NAME OF PROGRAM OFFICE REPRESENTATIVE Audrey Lee Silipo	MAIL CODE FF-S2-A	Gudrey Lee Silipo	DATE 4/14/98	
	year of the	X DISPOSITION W		
THIS FORM, WHEN COMPLETED, IS TO BE S	ENT TO YOUR	CENTER PUBLICATIONS OFFICE		

INSTRUCTIONS FOR COMPLETING THE NASA SCIENTIFIC AND TECHNICAL DOCUMENT AVAILABILITY AUTHORIZATION (DAA) FORM

Purpose. This DAA form is used to prescribe the availability and distribution of all NASA-generated and NASA-funded documents containing scientific and technical information (including those distributed via electronic media such as the World Wide Web and CD-ROMA).

Requirements. The author/originator must provide either a sultable summary description (title, abstract, etc.) or a completed copy of the document with this form. This form is initiated by the document author/originator and that individual is responsible for recommending/determining the availability/distribution of the document. The author/originator completes Sections I through III, and VI. The author/originator is also responsible for obtaining information and signature in Section IV to the extent the document discloses an invention for which patent protection has been applied. Subsequent to completion of these sections, the author/originator forwards the document to the appropriate Project Manager/Technical Monitor/Division Chief for further review and approval in Section VII, including a re-review of the planned availability and distribution. Once this approval is obtained, the DAA is forwarded to the NASA Headquarters or Center Export Administrator for completion of Section VIII. It is then forwarded for completion of Section IX to the cognizant NASA Headquarters Program Office or Delegated Authority, who provides final review and approval for release of the document as marked.

When to Use This Form. Documents containing STI and intended for presentation or publication (including via electronic media) must be approved in accordance with the NASA STI Procedures and Guidelines (NPG 2200.2). Documents that are to be published in the NASA STI Report Series must be coordinated with the appropriate NASA Headquarters or Center Scientific and Technical Information Office in accordance with NPG 2200.2. Note that information on the Report Documentation Page (if attached) is not to be entered on the DAA except for title, document date, and contract number.

How to Use this Form. Specific guidelines for each section of this form are detailed below.

- I. <u>Document/Project Identification</u>. Provide the information requested. If the document is classified, provide instead the security classification of the title and abstract. (Classified information must not be entered on this form). Include RTOP numbers on the Contract/Grant/Interagency/Project Number(s) line. Provide information on presentations or externally published documents as applicable.
- II. <u>Security Classification</u>. Enter the applicable security classification for the document, Documents, if classified, will be available only to appropriately cleared personnel having a "need to know."
- iii. Availability Category for Unclassified Documenta. Check the appropriate category or categories.

Export Controlled Document. If the document is subject to export restrictions (see NPG 2200.2, paragraph 4.5.3), the appropriate restriction must be checked, either International Traffic in Arms Regulations (ITAR) or Export Administration Regulations (EAR), and the appropriate United States Munitions List (USML) category or Commerce Control List (CCL), Export Control Classification Number (ECCN) must be cited.

Confidential Commercial Documents (Documents containing Trade Secrets, SBIR documents, and/or Copyrighted information). Check the applicable box (see NPG 2200.2 paragraph 4.5.7). When any of these boxes are checked, also indicate the appropriate limitation and expiration in the list to the right of these restrictions. These limitations refer to the user groups authorized to obtain the document. The limitations apply both to the initial distribution of the documents and the handling of requests for the documents. The limitations will appear on and apply to reproduced copies of the document. Documents limited to NASA personnel should not be made available to ensite contractors, if the Available Only With the Approval of Issuing Office limitation is checked, the NASA Center for AeroSpace Information will provide only bibliographic processing and no Initial distribution; CASI will refer all document requests to the Issuing office.

Publicity Available Document - Unrestricted Distribution. Check this box if the information in the document may be made available to the general public without restrictions (unrestricted domestic and international distribution). If the document is copyrighted (see paragraph 4.5.7.3 in NPG 2200.2), also check the "Copyrighted" box in this section.

- IV. <u>Document Disclosing an Invention</u>. This must be completed when the document contains information that discloses an invention (see NPG 2200.2, paragraph 4.5.9). When this box is checked, an additional appropriate availability category must be checked. Use of this category must be approved by NASA Headquarters or Center Patent Counsel or the Intellectual Property Counsel.
- V. <u>Stanket Release (Optional)</u>. Complete this optional section whenever subsequent documents produced under the contract, grant, or project are to be given the same distribution and/or availability as described in Sections II and III. More than one contract number or RTOP Number can be entered. This section may also be used to rescrind or modify an earlier Blanket Release. All blanket releases must be approved by the Program Office or its designee and concurred with by the Office of Management Systems and Facilities.
- VI. Author/Originator Verification. Required for all DAA forms.
- VII. <u>Project Officer/Technical Monitor/Division Chief Review.</u> The Project Officer/Technical Monitor/Author or Originator Division Chief or above must sign and date the form. The office code and typed name should be entered.
- VIII. Export Central Review/Confirmation. This section is to be completed by the authorized NASA Headquarters or Center Export Control Administrator for all documents.
- IX. Program Office or Delegated Authority Review. This section is to be completed by the duly authorized official representing the NASA Headquarters Program Office. Any delegation from NASA Headquarters to a NASA Center in accordance with NPG 2200.2 should be entered here.
- X. <u>Disposition</u>. For NASA Center use.

ESTIMATES OF PLANT CANOPY ATTENUATION, ABSORPTION AND REFLECTANCE SIGNATURES AND THEIR POTENTIAL TO ESTIMATE BIOPHYSICAL AND BIOCHEMICAL PLANT FEATURES

C. R. Hall^{1,2}, C. R. Bostater¹ and M. F. Pegaz^{1,2}

¹Marine & Environmental Optics Laboratory, Center for Remote Sensing Florida Institute of Technology, Melbourne, FL

²Dynamac Corporation, Kennedy Space Center, FL

ABSTRACT

Application of remote sensing technologies to ecosystem management requires a comprehensive understanding of radiative transfer theory and photon vegetation interactions at various spatial and temporal scales. As solar flux penetrates a plant canopy, absorption and scattering attenuate downward radiance and the light field diminishes. Biophysical and biochemical features of plant canopies that control attenuation include leaf area index, projected foliage coverage, leaf angle distribution, leaf moisture, chlorophyll, starch, cellulose, nitrogen and other organic and inorganic compounds. Field measurements of reflectance and bulk attenuation, transmission and absorption as a function of wavelength were made for canopies of live oak (Quercus viriginiana), Brazilian pepper (Schinus terebinthifolius) and grapefruit (Citrus paradisi) in east central Florida. Biophysical and biochemical analyses were conducted and Optimal Passive Ambient Correlation Spectroscopy (OPACS) was utilized to define optimal bands for use in predicting concentrations of biophysical and biochemical parameters.

1.0 INTRODUCTION

A primary objective for remote sensing in terrestrial ecology specifically and earth system science in general is determination of biochemical and biophysical features of vegetation, water, or land cover (Foody and Curran, 1994). Satellite and airborne remote sensing provide researchers access to spatial and temporal data necessary for

^{*} Presented at Eco-Informa '96, Lake Buena Vista, Florida, 4-7 November 1996

development, parameterization and testing of local, regional, and global scale ecosystem risk and environmental process models. Subtle changes in ecosystem functioning may be expressed in plant canopy biochemical or biophysical properties as a result of altered metabolic processes, carbon allocation patterns, and nutrient availability (Hoobs and Mooney, 1989). The potential to remotely estimate canopy constituents depends partly on the influence of the parameter on the reflectance signature. Potential applications of remote sensing techniques that allow for spatial and temporal estimation of biophysical and biochemical conditions include agricultural land management, exotic plant control, ecosystem risk modeling, environmental impact quantification and wildlife habitat management.

Peterson, et al., (1988) and others described attempts to remotely sense biochemical constituents of leaves and forest canopies utilizing the visible and infrared regions of the spectrum. These authors focused on stepwise linear regression techniques to describe relationships between leaf chemical concentrations and reflectance data. In the laboratory, using dried samples, these methods produced results that compared reasonably well with error rates for wet chemical analysis techniques. Statistically significant regressions were demonstrated for protein, starch, sugar, lignin, nitrogen, total chlorophyll and cellulose (Peterson, et al.1988).

However, Fourty et al., (1996) reported that the empirical relationships elaborated on for one site or sample had poor predictive performance when applied to other samples, sites or years. This lack of robustness and consistency drives the need to develop more analytical based approaches to describe the relationships between canopy reflectance and its biochemical and biophysical features that contribute to the high degree of variability observed in empirical results. These features include canopy structural parameters such as leaf angle, projected foliage coverage, leaf area index, canopy depth, bottom reflectance or background characteristics and scattering and absorption processes that take place at the leaf level. Analytical approaches to the solution of these problems must be based on radiative transfer theory and the associated solution and inversion of some form of the radiative transfer equations describing absorption, scattering and transmission features of the medium (Asar and Myneni, 1990; Fourty et al., 1996).

The objectives of this study were to evaluate potential uses of the optimal passive ambient correlation spectroscopy (OPACS) method developeed by Bostater, (1992) for analysis of high resolution spectral reflection, absorption and attenuation signatures of leaves and plant canopies. The OPACS procedure utilizes an optimal band selection methodology, based on a nonlinear second derivative estimator, to define spectral regions of absorption or backscatter that can be utilized in biophysical parameter estimation (Bostater et al., 1994a;1994b).

2.0 MATERIALS AND METHODS

Three plant canopies, live oak (Quercus viriginiana), Brazilian pepper (Schinus terebinthifolius) and grapefruit (Citrus paradisi) were chosen for this study based on differences canopy and leaf structure. Spectral measurements of downwelling and upwelling irradiance were made at the top, bottom, and at various depths within each of the canopies using an SE590 high radiometric and spectral resolution solid state spectrograph with a 252 channel linear diode array. Each channel samples approximately 3 nm of the electromagnetic spectrum over the spectral range from 368 to 1110 nm. Data are used to calculate reflectance, transmittance, absorptance and attenuation and bottom reflectance. Leaf samples were collected from each canopy, a video image of the leaves on a white background was captured for estimation of leaf area. The leaves were placed on ice in the dark, and taken to the laboratory for spectral analyses. An estimate of nadir viewing projected foliage coverage was derived by collecting ten random video images in the perpendicular plane and analyzing them for percent sky visible in the frame.

Leaf-level measurements of transmittance and reflectance were made in the Marine and Environmental Optics Laboratory at Florida Institute of Technology using a simultaneous reflectance and transmittancemeasuring system designed by Bostater (1994b). Sub-samples consisting of 10 leaves of each species were used for simultaneous measurements of leaf reflectance and transmittance. The thickness of each leaf was measured, to be used as a path length estimate for calculating leaf-level absorption coefficients. Leaf tissue was then freeze-dried, ground, and analyzed for chlorophyll, TKN and metals.

Software developed by Bostater, (1992) was utilized to develop estimates of optimal spectral bands for estimation of biophysical and biochemical parameters. The approach produces an estimate of the analytical solution to the second derivative of the complex radiative transfer equation for reflectance providing information on $d^2R/d^2\lambda$. (Bostater et al., 1994b). The second derivative inflection estimate is a precise measure of actual absorption or relative backscatter regions because it is based on three or more bands incorporating changes in spectral curvature.. The inflection estimator is defined as:

$$I(\lambda)_{\min} = R^{2}(\lambda)_{i} / [R(\lambda)_{i-n} \times R(\lambda)_{i+m}]$$
 '(1)

where $I_{(\lambda)}$ is the inflection or second derivative estimator centered at band i calculated from the reflectance signature. M and n are forward and backward operators, respectively (Grew, 1980). The optimal channel located at m, I, n is determined by computing all possible combinations of m, I, n ranging from i = 2-251, m = 2-252, and n = 1-250 (Bostater et al., 1994b).

3.0 RESULTS AND DISCUSSION

The objective of this study was to evaluate the potential for utilizing the OPACS software for definition of optimal bands useful in developing remote estimates of biophysical and biochemical characteristics of plants at the canopy and leaf scale. Canopy and leaf level measurements of reflectance, transmittance and absorptance were developed for the three species. Differences in spectral signatures are a result of variability in canopy structure, bottom reflectance, and absorption and backscatter of leaves and other plant structures. For example, Figure 1 presents canopy reflectance and transmittance signatures for citrus, brazilian pepper and live oak stands.




Figure 1. Example of reflectance and transmittance spectra for oak, citrus and brazilian pepper.collected at Kennedy Space Center in spring 1996. Note low values throughout photosynthetically active region between 400 - 700 nm.

Biophysical and biochemical measurements displayed subtle differences between the three canopies. Projected foliage coverage averaged 80% for citrus, 72% for oak and 66% for brazilian pepper. At a larger spatial scale citrus coverage would be lower than for the other species as a result of crown trimming and spacing between rows. This spacing issue may have contributed to the slight elevation observed in citrus transmittance estimates.

An example of OPAC analysis for total chlorophyll is presented in Fig. 2. Optimal bands were defined as 641, 697, and 1079 nm with a correlation of r = -0.81. The inflection point at 697 nm corresponds to the area of maximum chlorophyll absorption in the red region of the spectrum at the base of the red edge. Results of TKN analysis in general displayed weak correlation to the inflection estimates with most values below r = 0.70. The optimal bands were defined as 947, 1022, and 1088 nm with a correlation of r = 0.81.




Figure 2. Example of OPACS derived relationship between the inflection estimate and chlorophyll concentration for oak species on Kennedy Space Center, Fl. in spring 1995.

From results obtained in this study the optimum bands selected with OPACS define the inflection or spectral curvature occurring in association with biochemical and biophysical based absorption or backscatter regions. The unique nature of the OPACS procedure with it's strong basis in radiative transfer theory suggests that it is an optimal approach for quantitatively analyzing and interpreting high resolution spectral signatures. More detailed evaluations need to be conducted to address sources of variance in the original reflectance signal such as changing projected foliage coverage, seasonal variability in leaf chemistry and shape, and influences of bottom reflectance from leaf litter, ground cover and soils. Further testing and development in a larger variety of landscape types and situations, including application to high spectral resolution data derived from aircraft and satellites is needed to continue refining and testing the applicability of the OPACS procedure.

4.0 REFERENCES

Asar, G. and R. B. Myneni, "Application of Radiative Transfer Models for Remote Sensing of Vegetation Conditions", In Myneni, R. B. and J. Ross (Eds.) <u>Photon Vegetation Interactions</u>. Springer Verlag, New York. 565 p. 1990.

Bostater, C.R. "Remote sensing methods using aircraft and ships for estimating optimal bands and coefficients related to ecosystem responses". Proceeding of the First Thematic Conference on Remote Sensing for Marine and Coastal Environments. New Orleans, La. 1992.

Bostater, C. R., Ma, W-M., and A. Lamb. "Simulating radiative transfer in aquatic systems and contrasting results from ambient environmental spectroscopy: estuarine and near coastal waters." <u>Proceedings of the International Symposium on Spectral Sensing Research</u>. Gomez, R. B. (Ed). Volume II, 673-679. San Diego.1994a

Bostater, C. R., Hall, C. R., Vieglais, D., Rebbman, J., and M.J. Provancha. "Temporal measurements of high resolution spectral signatures of plants and relationships indicating water status." <u>Proceedings of the International Symposium on Spectral Sensing Research</u>. Gomez, R. B. (Ed). Volume I, 387-402. San Diego. 1994b.

Foody, G. and P. Curran. (Eds) Environmental remote sensing from regional to global scales. John Wiley & Sons. New York. 1994

Fourty, Th., F. Baret, S. Jacquemoud, G. Schmuck, and J Verbout. "Leaf optical properties with explicit description of its biochemical composition: direct and inverse problems." Remote Sens. Environ. 56:104-117.

Grew, G "Real-time test of MOCS algorithm during superflux 1980." Chesapeake Bay Plume Study, Superflux 1980, NASA Pub. 2188, Washington, D. C., 301-322. 1980.

Hoobs, R. J. and H. A. Mooney. (Eds) <u>Remote sensing of biosphere functioning</u>. Springer-Verlag. New York. 1989.

Peterson, D. L., Aber, J. D., Matson, P. A., Card, D. H., Swanberg, N., Wessman, C., and M. Spanner. "Remote sensing of forest canopy and leaf biochemical contents." Remote Sens. Environ. 24:85-108. 1988.